



Haddrell, A., & Thomas, R. (2017). Aerobiology: Experimental considerations, observations and future tools. *Applied and Environmental Microbiology*, 17(83).
<http://aem.asm.org/content/early/2017/06/26/AEM.00809-17?related-urls=yes&legid=aem;AEM.00809-17v1>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via the American Society for Microbiology at <https://doi.org/10.1128/AEM.00809-17> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

DSTL/JA100134

1 **Aerobiology: Experimental considerations, observations and future tools**

2 **Allen E. Haddrell¹ and Richard J. Thomas²**

3 *School of Chemistry, Cantock's Close, University of Bristol, BS8 1TS¹; Defence*

4 *Science and Technology Laboratory (Dstl), Porton Down, Salisbury, Wiltshire, UK,*

5 *SP4 0JQ²*

6

7 **Abstract**

8 Understanding airborne survival and decay of microorganisms is important for a
9 range of public health and biodefence applications including epidemiological and risk
10 analysis modelling. Techniques for experimental aerosol generation, retention in
11 aerosol phase and sampling require careful consideration and understanding so that
12 they are representative of the conditions the bioaerosol would experience in the
13 environment. This review explores current understanding of atmospheric transport in
14 relation to advances and limitations of aerosol generation, maintenance in the
15 aerosol phase and sampling techniques. Potential tools for the future are examined
16 at the interface between atmospheric chemistry, aerosol physics and molecular
17 microbiology that could explore heterogeneity and variability at the single droplet and
18 single microorganism level within a bioaerosol. The review highlights the importance
19 of method comparison and validation in bioaerosol research, and the benefits
20 application of novel techniques could bring to increased understanding of
21 aerobiological phenomena in diverse research fields, particularly during the
22 progression of atmospheric transport where complex interdependent
23 physicochemical and biological processes are occurring within bioaerosol particles.

DSTL/JA100134

24

25 **INTRODUCTION**

26 Aerosols injected into the atmosphere from the biosphere (bioaerosols) account for a
27 significant portion of all atmospheric aerosols (1). Despite their low numbers relative
28 to other natural aerosol, bioaerosols (whose sources include microorganisms
29 contained within windblown dust and sea spray) are speculated to impact climate
30 through behaving as efficient cloud condensation nuclei (2-3). Biological aerosols
31 are also important from the perspective of human health being intimately involved in
32 the transmission of many respiratory pathogens (4, 5).

33 Risk analysis modelling aims to develop predictive models of transmission and
34 infection based on laboratory generation of aerosols containing respiratory
35 pathogens. These experimental models are invaluable for understanding epidemic
36 transmission, developing infection control measures and advising bioterror
37 preparedness for public health (6-8). Effective risk modelling requires an in depth
38 understanding of experimental aerosol techniques and their potential impact on the
39 final outcome, whether that is aerosol decay, transmission rate or infectious dose.

40 This article reviews the current understanding, advances and limitations in laboratory
41 aerobiological studies where the relationship between microorganism preparation,
42 aerosol generation, evaporation, transport and fate cumulatively may affect the final
43 outcome of inhalational infection or survival in the environment. In this review, the
44 term "bioaerosol" will be limited to refer explicitly to infectious aerosol droplets
45 containing living species, specifically bacteria and viruses; the study of this subset of
46 bioaerosol comes with its own unique set of challenges that need to be recognized
47 and addressed. The PubMed database was searched to identify relevant studies

DSTL/JA100134

48 using the strings: aerosol AND survival, bioaerosol AND generation, bioaerosol AND
49 sampling. The terms bacteria and virus were interchanged for the term survival in the
50 first search string; only published studies were included. References with no relation
51 to bioaerosol as defined as 'infectious aerosol droplets' (e.g. fungal spores, pollen)
52 were generally discarded unless the technology could be applied to the field.
53 Retrieved studies were also reviewed for additional references. Although intrinsically
54 linked to the general theme of this review, the development of inhalational animal
55 models to replicate human disease is considered outside the scope and readers are
56 directed to an extensive literature in this field (9-11).

57

58 **AEROSOL GENERATION, SAMPLING AND POST-PROCESSING** 59 **CONSIDERATIONS**

60 Aerosol generation and sampling prior to microbiological analysis are conducted for
61 a range of bioaerosol related research activities (e.g. determination of aerosol decay
62 rates and inhalational infectious dose, efficacy of decontamination strategies, and
63 evaluation of bioaerosol sampling technologies). These dynamic processes can
64 cause damage due to shear forces acting on the microbial cells (12-27). Table 1
65 outlines some major aerosol generators and samplers used in aerobiological studies
66 and the operating mechanisms. The majority of studies use reflux aerosol generators
67 in conjunction with impingement to collect the generated aerosol. This system can be
68 safely used in biocontainment laboratories for inhalational challenges and aerosol
69 fate studies. However, comparative studies show that refluxing nebulizers produce
70 the greatest loss of physiological function as a function of time in bacteria (16, 19-21,
71 24). The loss of function has been linked to membrane damage (13, 20, 24), release

DSTL/JA100134

72 of ions into media (e.g. PO_4^{2-} ; 28), cell fragmentation (15, 23), reduction in ATP
73 activity (29) and magnitude of associated electrical charge (30) as the bacteria
74 remaining in the nebulizer repeatedly pass through the devices nozzle. Similar
75 effects are observed for viruses (25). Repair of bacterial cells damaged by
76 nebulization appears to be an energy dependent process with a requirement for
77 divalent cations although independent of *de novo* RNA or protein synthesis (13, 31);
78 it is unlikely that repair occurs in viruses due to their reliance on host cell factors for
79 protein transcription and translation. In contrast, it has been reported that damage is
80 reduced in non-refluxing aerosol generators where the microorganisms pass through
81 the nozzle once (16, 24).

82 Sampling methods for airborne microorganisms include impingement, impaction,
83 filtration, cyclonic separation, and electrostatic precipitation. This review will not
84 cover all bioaerosol samplers, rather selecting the main sampling mechanisms and
85 representative sampler models. The reader is directed to a couple of comprehensive
86 reviews on bioaerosol sampling for further detail (32, 33). Each sampling technique
87 has advantages and disadvantages for sampling microbial aerosols (Table 1) with
88 the potential to cause microbial damage. Dependent on the microbe this damage
89 may be transient: for example, impingement (AGI-30; 15 to 60 min) caused structural
90 damage to *Pseudomonas fluorescens* cells with recovery achieved on non-selective
91 media (15). Aerosol sampling times for determining infectious dose and aerosol
92 decay rates generally range from 1-10 min which minimize the effects of microbial
93 damage (22, 34). However, for infectious aerosols, few comparative studies of the
94 bioefficiency of different sampling mechanisms. Where studies comparing samplers
95 have been conducted, differences between microbial structures influence sampler
96 bioefficiency; for example, infectivity and culturability differences were observed

DSTL/JA100134

97 between bacteriophages and influenza A virions sampled by the SKC biosampler
98 and NIOSH cyclone (25, 35). Similar species dependent effects have been observed
99 for bacteria in sampling bioefficiency; in particular *Bacillus* endospores tend to be
100 less affected by aerosol sampling method (15, 17, 21, 22). One reason for
101 differences in sampler bioefficiency is variation in sampling velocities that for
102 impingement reaches $260 \text{ m}\cdot\text{s}^{-1}$, ten-fold greater than other samplers (36; Table 1).
103 Secondly, the rapid rehydration that occurs during sampling can be detrimental to
104 microorganisms (37-39).

105 Minimising stresses occurring during aerosol generation and sampling is hence
106 critical to accurate representation of aerosol decay and infectivity. Aerosol
107 generation stresses can be reduced by using single-pass devices that reduce the
108 probability of a microorganisms being damaged (24). Depending on sampler choice,
109 maximising recovery of microbes can be achieved in a number of ways. Prolonged
110 sampling times is a consistent cause of reduced viability and hence collection times
111 across all types of samplers and should be minimized (22, 40). The cell membrane is
112 a major site of damage for Gram negative bacteria being aerosolised as sampled,
113 demonstrated by increased sensitivity to hydrolytic enzymes (12). Impingement
114 requires collection into liquid which can be optimised to reduce osmotic shock and
115 maximise repair and recovery. For example, addition of compatible solutes and
116 scavenging enzymes (i.e. trehalose, raffinose, polyhydric alcohols, betaine and
117 catalase) can facilitate survival following the stresses associated with aerosol
118 generation, transport and sampling (38, 41-46). Particle bounce and viability loss in
119 impactors for vegetative *Bacillus subtilis* and *Escherichia coli* cells was reduced by
120 applying a thin film of mineral oil significantly enhancing collection efficiency (47).
121 Filtration methods provide high physical collection efficiencies, but bioefficiency can

DSTL/JA100134

122 be dependent on filtration time and post-processing procedures (21, 24, 48, 49). A
123 major problem with filtration samplers is continued drawing of air through the filter
124 desiccates collected microorganisms in a time-dependent manner. However,
125 filtration onto gelatin membranes provides a medium that retains moisture and can
126 be placed into warm media to recover collected microorganisms providing good
127 bioefficiency (21, 24).

128 Post-sampling enumeration and storage are additional considerations. Enumeration
129 can introduce error as organisms can be sensitive to impaction onto an agar surface
130 (50), sensitive to the plating media (15) and the process of spread plating (51-53).
131 Direct methods such as microscopy or flow cytometry in conjunction with various
132 dyes or quantitative polymerase chain reaction (PCR) can indicate physiological
133 activity of the collected microorganisms (15, 17, 54). Storage temperature, sampling
134 solution and length of time can prompt microbial replication (or death) causing
135 misrepresentation of the actual viability of the sampled bioaerosol (48). Samples
136 should be processed as soon as possible after aerosol sampling; however this is
137 highly dependent on the microorganism as for example, *Bacillus* endospores have
138 been demonstrated to be less affected by storage temperature (4 and 25 °C)
139 compared to *Escherichia coli*; however compared to immediate enumeration, both
140 species had increased counts after extended periods of storage at 25 °C (10 and 24
141 h for *B. subtilis* and *E. coli* respectively) indicating significant disaggregation and/or
142 multiplication in the collection medium, which in this case was sterile deionized water
143 containing a small quantity of detergent (48).

144 The data indicates that the method of aerosol generation can damage the
145 microorganism at the subcellular level, at the very least subtly, and influence
146 resultant estimates of microbial viability in the aerosol phase. None of these

DSTL/JA100134

147 mechanisms are entirely representative of the natural transmission mechanisms of
148 respiratory pathogens e.g. coughing and sneezing followed by deposition in the
149 respiratory tract (4, 5). The complexity of fluid fragmentation and droplet formation of
150 oro-respiratory secretions during coughs and sneezes has recently been elucidated
151 with the viscoelastic properties of respiratory secretions playing a defining role in
152 final droplet size (55, 56). Viscoelasticity of respiratory secretions will change with
153 anatomical location (e.g. nasal, bronchial) and disease state (e.g. chronic bronchitis,
154 sinusitis, cystic fibrosis) as a result of changes in mucin content which will also affect
155 droplet sizes (57, 58). Natural aerosol transmission events are likely to be less
156 violent than the aforementioned aerosol generation processes. Therefore, selection
157 and validation of experimental regimes (aerosol generator, spray fluid composition
158 and sampling) to minimize microbial damage, promote maximal recovery and most
159 closely replicate the natural event being modelled, is important for interpretation of
160 aerosol data used in risk analysis models. Based on this review, and more extensive
161 reviews on sampling methodology (32, 33) it is apparent that given the variability in
162 microorganisms responses to the stresses of aerosol generation and collection, then
163 it is advisable to perform method validation for each particular microorganism.
164 Testing a range of aerosol generators and samplers to ensure the behaviour of the
165 microorganism within the system is understood facilitates appropriate selection of
166 apparatus and methodology to maximise recovery during enumeration.

167 **AEROSOL TRANSPORT AND PHYSICAL PROCESSING**

168 The physicochemical properties of bioaerosol particles govern all of the biological
169 processes within. The conditions in a bioaerosol particle that a microorganism will
170 experience can be dramatically different than in bulk liquid; the solute concentrations
171 commonly reach supersaturation (59), while the rate of water transport within the

DSTL/JA100134

172 droplet can vary by orders of magnitude (60). Both of these properties are regulated
173 by total water present in the droplet. Thus a detailed understanding of the
174 hygroscopic properties of a bioaerosol as a function of solute composition (including
175 biological species itself) is critical for understanding and predicting longevity and
176 overall infectivity.

177 The typical trajectory in RH for a respiratory pathogen would be high at the point of
178 dispersion (>95%) to low during atmospheric transport (ambient relative humidity,
179 RH) to high upon inhalation (>95%) (61). During its lifetime, the water activity (a_w)
180 within a droplet equilibrates with the atmospheric RH through either the addition or
181 removal of water (62). From droplets larger than 100 nm in size, the water activity is
182 equal to the gas phase RH at equilibrium. The rate at which this mass flux occurs
183 and the final particle size attained are a reflection of the temperature and humidity of
184 the gas phase of the aerosol and the droplet solute (63, 64). Importantly, all
185 microorganisms require water for activity as critical enzyme driven biochemical
186 reactions (e.g. respiration). Interestingly, in studies looking at osmotic tolerance in
187 bulk liquid phase, depending on bacterial species, multiplication and growth is
188 inhibited at a_w values of 0.86 – 0.97 with further reductions inducing dormancy or
189 eventually reducing viability (65, 66).

190 The hygroscopic behaviour of any multicomponent aerosol is dependent on the
191 relative abundance of each chemical species in the solute, where each component
192 will contribute a proportion to the uptake or loss of water (62). This paradigm holds
193 true for bioaerosol, for example it has been shown that the solute concentration
194 affects the hygroscopic growth of aerosolized *B. subtilis* and *Pseudomonas*
195 *fluorescens* vegetative cells (67). However, to study the hygroscopic behaviour of
196 aerosol where the aim is to generate predictive models, much information about the

DSTL/JA100134

197 solute is required. The relative abundance of each component within the aerosol is
198 mandatory (68-72), as is a detailed understanding of how the various components
199 within the solute interact with one another (73). While this is somewhat straight
200 forward with regards to non-biological aerosol, it remains a major challenge in
201 bioaerosols. For example, infected individuals coughing and sneezing will produce
202 larger droplets with different concentrations of mucus and other organic and
203 inorganic solutes compared to healthy individuals (58). Similarly, in laboratory
204 studies, microbial culture conditions (liquid broth, solid agar and nutrient
205 composition) and growth phase affect the concentration and types of nutrients
206 present in the spray suspension and these factors influence aerosol survival (25, 74-
207 78). Indeed, survival of a viral simulant, the bacteriophage MS2, differed in human
208 derived saliva, artificial saliva and cell culture medium, with greatest decay observed
209 in human derived saliva (79). This has been observed for other viruses and bacteria
210 upon comparing survival after aerosolization from body fluids (natural or synthetic)
211 and culture medium (80-83). This highlights the caution needed in extrapolation of
212 results from the experimental to *in vivo* situations being modelled in risk analysis.

213 The primary challenge in experimental studies of the factors that regulate the
214 hygroscopic behaviour of bioaerosol is to control and know the complete composition
215 of the bioaerosol droplets. For example, a simple factor such as control of the
216 number of organisms per droplet/particle is not trivial using conventional
217 aerosolization processes. To attempt to address this specific issue in studies of
218 laboratory generated bioaerosols, a particular size is selected for a nebulized and
219 dried bioaerosol sample allowing estimation of the number of species per droplet
220 prior to hygroscopic analysis (16). For more complex (and atmospherically relevant)
221 bioaerosol, the hygroscopic behaviour of anthropogenic bioaerosol has been

DSTL/JA100134

222 estimated indirectly (84, 85). In these studies, the relative growth in bioaerosol
223 particle size with increases in RH was estimated through correlation analysis
224 between the temporal size distributions (aerodynamic diameter) of airborne fungi
225 with meteorological information (RH).

226 Thermodynamic models to predict the hygroscopic behaviour of aerosol (e.g.
227 Universal Quasichemical Functional Group Activity Coefficients; UNIFAC) have been
228 used for bioaerosols to limited success (59, 86). Generally, these models are able to
229 predict the hygroscopic behaviour of large and complex organic molecules through
230 parameterization of the functional groups present (such as carboxylic acids; 87).
231 Even though, organically, bioaerosol consists primarily of sugar alcohols and highly
232 polar sugars (88), it remains unclear the extent to which these models can be used
233 to predict the hygroscopic behaviour of bioaerosols (89). The reason for this is that
234 even when the relative abundances of functional groups and chemical species within
235 a single bioaerosol droplet are known, the accumulation of noncovalent interactions
236 between these species is not; the presence of cellular membranes within the droplet
237 could kinetically limit the hygroscopic behaviour of all the chemical species within the
238 aerosol.

239 The limited number of comprehensive studies that explicitly study the
240 physicochemical properties of bioaerosol is problematic. Their absence has
241 constrained the means by which the longevity of suspended bioaerosol can be
242 investigated.

243

244 **DETERMINING BIOAEROSOL LONGEVITY**

DSTL/JA100134

245 Bioaerosol longevity is simply the length of time in which a biological species will
246 remain either infectious or viable while suspended as a single particle. In an ideal
247 experiment, the entire composition of the target bioaerosols would be explored; as
248 discussed in previously sections this is technically challenging due to the selectivity
249 of samplers and the heterogeneity of bioaerosol composition. Despite this, numerous
250 studies on bioaerosol longevity have been published.

251 Techniques for investigating survival of bioaerosols *in vitro* (Table 2) tend to either
252 maintain the particles in the air column (i.e. 'dynamic bioaerosols') or captured on
253 fine substrate such as spider silk or glue fibres (i.e. 'captured bioaerosols'). The
254 rotating drum is probably the standard procedure used for aerosol longevity studies
255 based on Goldberg and colleagues seminal design (90). Modifications have
256 permitted greater control (e.g. *in situ* monitoring of parameters) and accessibility to a
257 range of environmental parameters (e.g. temperature, UV, volatile organic
258 compounds), and the suspension of larger aerosol particle sizes for sufficiently long
259 periods of time (91-94). Methods based on capturing bioaerosols on microfibers
260 derived from spider escape silk and glue gun fibres have been utilised with success
261 (78, 95-97). Comparative studies on filoviruses have demonstrated that microthread
262 captured bioaerosols decay at a similar rate as those held dynamically within rotating
263 vessels (34, 98).

264 The methods for retention of microorganisms in the aerosol phase have been used
265 extensively to determine biological decay in the airborne state as a function of time
266 and a range of environmental conditions (Table 3). The aerosol is sampled at time
267 intervals and the number of viable microorganisms enumerated enabling calculation
268 of aerosol decay rate. Sampling method and subsequent microbiological processing
269 and enumeration can alter the number of recovered microorganisms (15, 17, 21, 22).

DSTL/JA100134

270 Therefore it is important to minimize microbial stress during aerosol collection to
271 facilitate accurate calculation of the decay rate. During method validation, it is
272 important to differentiate biological decay from physical losses due to deposition on
273 the walls of the vessel or removal from the microthreads due to turbulence (or the
274 presence of antimicrobial substances on the silk). Physical loss in aerosol systems is
275 determined by using physical tracers that will not biologically decay such as *Bacillus*
276 spores, chemicals (e.g. fluorescein) or polymer beads (21, 99, 100). The decay rates
277 of the target microorganism and the physical tracer can be compared and the true
278 biological decay rate determined.

279 A disadvantage of these techniques is that they sample bulk aerosol and it is difficult
280 to develop an appreciation of microenvironment heterogeneity occurring within
281 individual aerosol droplets from the physicochemical and biological perspective. For
282 example, each individual aerosol droplet is likely to have a different chemical
283 composition, exacerbated by differences in particle size that manifest themselves
284 biologically on the microorganisms incorporated within the droplets. Such differences
285 may be a source of variability in how microbes respond and survive aerosol
286 transport.

287

288 ENVIRONMENTAL FACTORS AFFECTING MICROBIAL LONGEVITY DURING 289 ATMOSPHERIC TRANSPORT AND BACTERIAL SURVIVAL MECHANISMS

290 A large number of environmental and meteorological factors can influence microbial
291 survival during aerosol transport (Table 3), and to provide greater context for
292 interpretation of results the environmental features of the sampling site should be
293 described. The fate of the microorganism is likely dictated by its physiological status

DSTL/JA100134

294 which is a combinatorial consequence of the atomisation process (e.g. spray device,
295 cough, sneeze) with the associated evaporative stresses of aerosol transport and
296 rehydration during inhalation (or sampling into liquid). The mechanisms by which the
297 microorganisms perish have been partially elucidated and depend on the
298 composition of the droplet and surrounding atmosphere.

299 Atmospheric oxidants (e.g. reactive oxygen and nitrogen species, sulphur dioxide,
300 ozone) will impact on microbial longevity by either directly acting on the organism or
301 with constituents within the aerosol droplet (101, 102). Presence of oxygen has been
302 demonstrated to have a deleterious effect on airborne coliform bacteria, particularly
303 at RH less than 40%, and hypothesised to be due to production of reactive oxygen
304 species (ROS) by Maillard reactions (31, 103). Maillard reactions are amino-carbonyl
305 reactions occurring between amino groups on proteins and reducing sugars that
306 cause oxidation of macromolecules and death in microorganisms (104). In airborne
307 microorganisms, these reactions may be the cause of oxidative damage to critical
308 enzymes (44, 105-107), phospholipids and nucleic acids causing metabolic
309 imbalance, destabilisation of membranes and reducing repair activity (31).
310 Interestingly, recently Maillard chemistry has been implicated as a source of organic
311 compounds within atmospheric aerosols altering particle viscosity and hence the
312 diffusivity rate of water and reactive gases (108). Bioaerosols (including virus,
313 vegetative bacteria, spores and peptides) subjected to atmospheric ozone
314 concentrations and variations in RH showed temporal changes in fluorescence
315 spectra related to oxidation and hydrolysis of tryptophan (109-111). Although
316 survival is generally greater at higher RH (>80%), certain values (i.e. 70-85% RH for
317 *E. coli* B; 41, 44) produce a large decrease in aerosol survival (41, 107, 112, 113).
318 Likewise, RH dependent changes in salt concentrations and pH within droplets

DSTL/JA100134

319 influence virus viability causing conformational changes in surface proteins and
320 membrane fluidity affecting infectivity (114).

321 Solar irradiation and atmospheric pollutant gases (including open air factor; OAF) are
322 two further environmental parameters that can significantly affect longevity in the
323 aerosol phase. Solar irradiation markedly decreased viability compared to control
324 conditions that simulate the night (46, 78, 115-118). Particle size-dependent survival
325 against solar irradiation has been observed with bacterial clusters persisting for
326 longer periods (78, 117). Terrestrial solar spectral irradiance varies through the day,
327 with season and with geographical location (119). The UV wavelengths are of most
328 importance for inactivating microorganisms (116, 117), where UV-A and UV-B reach
329 the troposphere with the potential to cause a variety of DNA genomic lesions and
330 damage to nucleic acids, proteins and lipids due to generation of reactive oxygen
331 species (120-121). It is important that studies using both simulated and natural solar
332 irradiation report variables such as solar intensity as accurately as is reasonably
333 possible to facilitate data interpretation and standardisation between laboratories.

334 Atmospheric constituents such as various pollutant gases and secondary organic
335 aerosols (SOAs; Table 3) have been demonstrated to have significant deleterious
336 effects on aerosol longevity (31, 93, 122, 123-130). Many of these may contribute to
337 a phenomenon known as 'open air factor' (OAF) where aerosolized microorganisms
338 exposed to open climatic conditions decay more rapidly than those in enclosed
339 laboratory vessels subjected to similar temperature and RH (31, 123-125, 129, 130).
340 The precise nature of OAF is not fully understood but is hypothesised to involve a
341 number of highly reactive products (e.g. hydroxyl radicals) from photochemical
342 interactions between ozone and unsaturated hydrocarbons from anthropogenic (e.g.
343 engine-related alkenes) and non-anthropogenic sources (e.g. plant terpenes) (31,

DSTL/JA100134

123). The reactive species rapidly oxidise and degrade macromolecules such as lipids, proteins and nucleic acids (31, 131). The effect of OAF is enhanced at high humidity (80-90% RH) for both *E. coli* and *Micrococcus albus* (123). Such humidity effects warrant investigation, possibly relating to the increased water content of aerosol particles at higher humidity.

How microbes regulate and survive aerosol transport is undetermined. Evidence suggests that the ability for transcription and translation to occur in the environment of an evaporating droplet is reduced (31, 132, 133). Evaporation and rehydration of aerosol particles imparts osmotic and desiccative stress on the microbe reflective of the humidity of the surrounding atmosphere and composition of the particle. The molecular response of many bacterial species to osmotic stress and desiccation is well documented from research understanding survival in food matrices, aquatic and marine systems and terrestrial environments (66). Hyperosmotic stress (i.e. increased a_w) causes a reduction in cytoplasmic volume as water exits the bacterium; concomitantly cell growth and respiration cease as the bacterium adapts to the hyperosmotic conditions. Initially charged solutes (e.g. K^+ ions, glutamate) are accumulated via specific uptake mechanisms (66, 134-136). Interestingly, inability to control efflux of K^+ ions correlated with decreased survival in aerosolised *E. coli* cells (28, 137). Synthesis of compatible solutes (e.g. trehalose) or uptake from the surrounding media (e.g. glycine betaine, proline) stabilises proteins, enzymes and membrane phospholipids enable critical biochemical processes to continue in hyperosmotic stressed bacteria. As the bacterial cell stabilises, a number of proteins are synthesised prompting repair of DNA damage, scavenging of reactive oxygen species and degradation of misfolded proteins (66, 134-136). Osmotically adapted cells often show cross-tolerance to other stresses such as high temperature and

DSTL/JA100134

369 oxidative shock (138). Recently, *E. coli* subjected to rapid downshift in a_w (0.993 to
370 0.960) in media was demonstrated to control protein misfolding by transient
371 expression of the RpoE and RpoH regulons in conjunction with the RpoS regulon to
372 facilitate prolonged adaptation to the hyperosmotic conditions (139).

373 The molecular studies described above have all been conducted in bulk solution
374 phase and expose the microorganisms to hyperosmotic stress. Microorganisms will
375 be exposed to hyperosmotic conditions within an evaporating droplet (i.e. low a_w
376 conditions), enabling speculation that similar molecular mechanisms play role in
377 bacterial survival within evaporating aerosol droplets. As will be discussed later,
378 advances in atmospheric chemistry and single cell genomic techniques means that
379 investigation of whether similar molecular mechanisms occur in an aerosol droplet as
380 a function of evaporation rate and droplet composition are on the horizon.
381 Importantly, if airborne microorganisms can induce adaptive responses promoting
382 survival then there is the potential that colonisation and infection of the respiratory
383 tract is primed whilst the bacteria are transported in the atmosphere. Any induced
384 virulence factors would offer attractive targets for combating respiratory infection.

385

386 **NEW TECHNIQUES FOR ADVANCING AEROSOL SCIENCE AND** 387 **AEROBIOLOGY**

388 Bioaerosols, even when produced under controlled laboratory conditions, are
389 complex. They are generally polydisperse in terms of both physicochemical and
390 biological properties, and the heterogeneity in the nature of the bioaerosol evolves
391 with time and distance from the source. Technological advances in the fields of
392 aerosol science and molecular biology are timely to facilitate multidisciplinary

DSTL/JA100134

393 approaches to understand heterogeneity at the single droplet and single
394 microorganism level (including microbial aggregates) and to explore the
395 fundamentals of biological decay and survival in aerosol droplets.

396 Optical techniques such as optical tweezers and electrodynamic balances where
397 single aerosol droplets can be captured and levitated within an electric field for
398 periods of time (seconds to days) have been extensively used in atmospheric
399 chemistry to investigate heterogeneous chemistry, phase separation, hygroscopicity
400 and ice nucleation activity using analytical techniques including Raman
401 microspectroscopy (140-145). Utilisation of these techniques for biological aerosol
402 has been limited to date. However, optically trapped single biological cells in solution
403 produce characteristic Raman scattering signatures (146-149) and *E. coli* exposed to
404 1-butanol resulted in spectroscopic and anisotropic detection of real-time phenotypic
405 changes in fatty acid composition and membrane fluidity (149). Although these
406 studies were conducted in liquid bulk solution rather than aerosol droplets, it
407 exemplifies the power of the technology. Furthermore, such techniques are being
408 used to explore individual aerosol particles containing microorganisms, fungal spores
409 and pollen (150-152). The electrodynamic balance technique has been used to
410 accurately deposit single particles containing respiratory syncytial virus onto airway
411 epithelial cells enabling the cellular response to infection to be analysed (153). This
412 technique enables interaction at the air-cell interface with single aerosol particles, a
413 more representative scenario than the air-liquid interface studies commonly
414 conducted for *in vitro* infection studies. It is a technique that seems applicable
415 although currently rarely applied to understanding the heterogeneity of bioaerosols at
416 the single droplet and microorganism level.

DSTL/JA100134

417 Microbial cells respond to environmental stimuli by regulating gene expression
418 resulting in modulation of the quantities and composition of functional proteins
419 available to combat a particular stressful condition. Transcriptional analysis and
420 insertional mutagenesis have been used to identify bacterial genes regulated in
421 response to stresses associated with aerosol survival such as desiccation, and
422 osmotic pressure (136, 154). Currently, these techniques have not been applied to
423 aerosolised microbial populations, however it can be hypothesised that similar
424 responses may be expected and warrant exploration. The relative abundance of
425 particular proteins critical to aerosol survival will vary from cell to cell. Exploring this
426 heterogeneity at the single cell level is complicated due to the relatively low
427 abundance of stress-responsive proteins. However, the last five years have seen
428 significant advances in molecular techniques enabling exploration of the genomic,
429 proteomic or 155-158). Techniques for isolating single cells such as flow cytometry
430 and microfluidics can be combined with techniques such as PCR and next-
431 generation sequencing for probing the transcriptional response of single cells (159).
432 Indeed, single cell genomic techniques have been applied to understanding airborne
433 metagenomes in urban settings (160, 161). Application to aerosolised populations in
434 a laboratory setting would seem straightforward. However, care in experimental
435 design would be needed to discriminate the true effects of aerosol transport and the
436 stresses of aerosol generation and sampling.

437 These emerging technologies have the potential to dramatically impact numerous
438 areas of bioaerosol science. They will lead to improved parameterization of the
439 fundamental properties of bioaerosol, such as the interplay between environmental
440 conditions with species longevity and/or gene expression. This data will lead to
441 better predictions of disease dynamics in areas such as general industrial hygiene,

DSTL/JA100134

442 animal husbandry, hospital design and biosecurity. Furthermore, the data collected
443 from these laboratory based instruments will inform conventional research of
444 environmental samples.

445

446 **CONCLUDING REMARKS**

447 Experimental factors affect the microbiological sample taken forward for
448 quantification of infectious dose or biological decay rate. Therefore a thorough
449 understanding of the sampling and enumeration process is critical to interpretation of
450 the final data set. Furthermore, no single aerosol generation or sampling method is
451 likely to suit all purposes (i.e. size selectivity, species sensitivity), therefore the
452 experimental apparatus should be selected based on the hypothesis and
453 microorganism being tested and the data interpreted alongside the caveats
454 associated with the methodology. For experiments designed to generate data for
455 input into risk analysis determination of human inhalational exposure then it is
456 recommended that aerosol generators, samplers (and collection fluid) be used that
457 cause minimal damage or promote maximal recovery of the microorganisms during
458 collection to prevent underestimation of risk estimates.

459 Fundamental questions remain regarding aerosol transmission of respiratory
460 pathogens, particularly the underlying mechanisms of survival and/or death during
461 aerosol transport and the role the microenvironment of the droplet plays as it
462 evaporates then rehydrates during inhalation. However, as outlined in this review,
463 advances in distinct scientific fields could support a systematic dissection of the
464 biological response of microorganisms within compositionally controlled aerosol
465 droplets within specific atmospheric conditions. It is envisaged that within the next

DSTL/JA100134

466 ten years multidisciplinary approaches combining existing and novel techniques in
467 atmospheric chemistry, aerobiology and molecular biology will converge and begin to
468 dissect and empirically understand the mechanisms of microorganisms survival and
469 decay in the aerosol state and the effect on infectivity and disease transmission.

DSTL/JA100134

470 **ACKNOWLEDGEMENTS**

471 Dr Allen Haddrell and Dr Richard Thomas recognise the funding support of Elizabeth
472 Blackwell Institute for Health Research and the Ministry of Defence respectively. Thanks
473 are extended to Prof Jonathon Reid for critical review of the manuscript.

474

475 **REFERENCES**

- 476 1. Jaenicke, R. 2005. Abundance of cellular material and proteins in the atmosphere.
477 *Science* 308:73-73.
- 478 2. Sesartic A, Lohmann U, Storelvmo T. 2013. Modelling the impact of fungal spore ice
479 nuclei on clouds and precipitation. *Environ Res Lett* 8:doi:10.1088/1748-
480 9326/8/1/014029.
- 481 3. Konstantinidis KT. 2014. Do airborne microbes matter for atmospheric chemistry and
482 cloud formation? *Environ Microbiol* 16:1482-1484.
- 483 4. Gralton J, Tovey E, McLaws M-L, Rawlinson WD. 2011. The role of particle size in
484 aerosolised pathogen transmission: A review. *J Infect* 62:1-13.
- 485 5. Fernstrom A, Goldblatt M. 2013. Aerobiology and its role in transmission of infectious
486 diseases. *J Pathog* 2013:1-13. <http://dx.doi.org/10.1155/2013/493960>.
- 487 6. Lander W, van Kerckhove K, Chao DL, Hens N, Beutels P. 2012. A nice day for an
488 infection? Weather conditions and social contact patterns relevant to influenza
489 transmission. *PLoS One* 7:e48695.
- 490 7. Hong T, Gurian PL, Huang Y, Haas CN. 2012. Prioritizing risks and uncertainties
491 from intentional release of selected Category A pathogens. *PLoS One* 7:e32732.

DSTL/JA100134

- 492 8. Milton DK. 2012. What was the primary mode of smallpox transmission? Implications
493 for biodefense. *Front Cell Infect Microbiol* 2:1-7.
- 494 9. Kukavica-Ij I, Levesque RC. 2008. Animal models of chronic lung infection in
495 *Pseudomonas aeruginosa*: useful tools for cystic fibrosis studies. *Lab Anim* 42:389-
496 412.
- 497 10. Thangavel RR, Bouvier NM. 2004. Animal models for influenza virus pathogenesis,
498 transmission and immunology. *J Immunol Meth* 410:60-79.
- 499 11. Swearingen JR. 2012. *In Biodefense Research Methodology and Animal Models*.
500 CRC Press.
- 501 12. Hambleton P. 1970. The sensitivity of Gram-negative bacteria, recovered from
502 aerosols, to lysozyme and other hydrolytic enzymes. *J Gen Microbiol* 61:197-204.
- 503 13. Hambleton P. 1971. Repair of wall damage in *Escherichia coli* recovered from an
504 aerosol. *J Gen Microbiol* 69:81-88.
- 505 14. Kang Y-J, Frank JF. 1989. Biological aerosols: a review of airborne contamination
506 and its measurement in dairy processing plants. *J Food Prot* 52:512-524.
- 507 15. Terzieva S, Donnelly J, Ulevicius V, Grinshpun S, Willeke K, Stelma GN & Brenner
508 KP. 1996. Comparison of methods for detection and enumeration of airborne
509 microorganisms collected by liquid impingement. *Appl Environ Microbiol* 62:2264-
510 2272.
- 511 16. Reponen T, Willeke K, Ulevicius V, Grinshpun SA, Donnelly J. 1997. Techniques for
512 dispersion of microorganisms into air. *Aero Sci Technol* 27:405-421.
- 513 17. Rule AM, Kesavan J, Schwab KJ, Buckley TJ. 2007. Application of flow cytometry for
514 the assessment of preservation and recovery efficiency of bioaerosols samplers
515 spiked with *Pantoea agglomerans*. *Environ Sci Technol* 41:2467-2472.

DSTL/JA100134

- 516 18. Hermann JR, Zimmerman JJ. 2008. Analytical sensitivity of air samplers based on
517 uniform point-source exposure to airborne Porcine reproductive and respiratory
518 syndrome virus and swine influenza virus. *Can J Vet Res* 72:440-443.
- 519 19. Eninger RM, Hogan CJ, Biswas P, Adhikari A, Reponen T, Grinshpun SA. 2009.
520 Electro spray versus nebulization for aerosolization and filter testing with
521 bacteriophage particles. *Aero Sci Technol* 43:298-304.
- 522 20. Thomas RJ, Webber D, Hopkins R, Frost A, Laws T, Jayasekera P, Atkins T. 2011.
523 The cell membrane as a major site of damage during aerosolization of *Escherichia*
524 *coli*. *Appl Environ Microbiol* 77:920-925.
- 525 21. Dabisch P, Yeager J, Kline J, Klinedinst K, Welsch A, Pitt ML. 2012a. Comparison of
526 the efficiency of sampling devices for aerosolized *Burkholderia pseudomallei*. *Inhal*
527 *Toxicol* 24:247-254.
- 528 22. Dabisch P, Yeager J, Kline J, Klinedinst K, Welsch A, Pitt ML. 2012b. Recovery
529 efficiencies for *Burkholderia thailandensis* from various aerosol sampling media.
530 *Front Cell Infect Microbiol* 2:1-7.
- 531 23. Zhen H, Han T, Fennell DE, Mainelis G. 2013. Release of free DNA by membrane-
532 impaired bacterial aerosols due to aerosolization and air sampling. *Appl Environ*
533 *Microbiol* 79:7780-7789.
- 534 24. Zhen H, Han T, Fennell DE, Mainelis G. 2014. A systemic comparison of four
535 bioaerosols generators: affect on culturability and cell membrane integrity when
536 aerosolizing *Escherichia coli* bacteria. *J Aero Sci* 70:67-79.
- 537 25. Turgeon N, Toulouse M-J, Martel B, Moineau S, Duchaine C. 2014. Comparison of
538 five bacteriophages as models for viral aerosol studies. *Appl Environ Microbiol*
539 80:4242-4250.

DSTL/JA100134

- 540 26. Fennelly KP, Tribby MD, Wu CY, Weil GL, Radonovich LJ, Loeb JC, Lednicky JA.
541 2015. Collection and measurement of aerosols of viable influenza virus in liquid
542 media in an Andersen cascade impactor. *Virus Adapt Treat* 7:1-9.
- 543 27. Han T, Wren M, DuBois K, Therkom J, Mainelis G. 2015. Application of ATP-based
544 bioluminescence for bioaerosol quantification: effect of sampling method. *J Aerosol*
545 *Sci* 90:114-123.
- 546 28. Anderson JD, Dark FA. 1967. Effects of aerosolization upon survival and potassium
547 retention by various bacteria. *J Gen Microbiol* 52:99-105.
- 548 29. Han T, Wren M, DuBois K, Therkom J, Mainelis G. 2015. Development of ATP
549 Bioluminescence method for rapid bioaerosol quantification. *J Aero Sci* 90:114-123,
- 550 30. Mainelis G, Willeke K, Baron P, Reponen T, Grinshpun SA, Gorny R, Trakumas S.
551 2001. Electrical charges on airborne microorganisms. *Aero Sci* 32:1087-1110.
- 552 31. Cox CS. 1987. The aerobiological pathway of microorganisms. John-Wiley & Sons,
553 New York.
- 554 32. Kesavan J, Sagripanti J-L. 2015. Evaluation criteria for bioaerosol samplers. *Environ*
555 *Sci Process Impact* doi:10.1039/c4em00510d
- 556 33. Grinshpun S, Buttner M, Mainelis G, Willeke K. 2016. Sampling for airborne
557 microorganisms. In Yates, M, Nakatsu, C, Miller, R, Pillai, S (ed), *Manual of*
558 *Environmental Microbiology*, 4th ed. ASM Press, Washington, DC
- 559 34. Piercy TJ, Smither SJ, Steward JA, Eastaugh L, Lever MS. 2010. The survival of
560 filoviruses in liquids, on solid substrates and in a dynamic aerosol. *Appl Microbiol*
561 109:1531-1539.

DSTL/JA100134

- 562 35. Cao G, Noti JD, Blachere FM, Lindsley WG, Beezhold DH. 2011. Development of an
563 improved methodology to detect infectious airborne influenza virus using the NIOSH
564 bioaerosol sampler. *J Environ Monit* 13:3321-3328.
- 565 36. Mainelis G, Grinshpun SA, Willeke K, Reponen T, Ulevicius V, Hintz PJ. 1999.
566 Collection of airborne microorganisms by electrostatic precipitation. *Aero Sci*
567 *Technol* 30:127-144.
- 568 37. Benbough JE. 1971. Some factors affecting the survival of airborne viruses. *J Gen*
569 *Virol* 10:209-220.
- 570 38. Marthi B. 1994. Resuscitation of microbial aerosols. p 192-225. *In* Lighthart B, Mohr
571 AJ (eds), *Atmospheric microbial aerosols: theory and applications*. Springer Science,
572 New York.
- 573 39. Lang E, Zoz F, Iaconelli C, Guyot S, Alvarez-Martin P, Beney L, Perrier-Cornet J-M,
574 Gervais P. 2016. Recovery estimation of dried foodborne pathogens is directly
575 related to rehydration kinetics. *PLoS One* 11:e0160844.
- 576 40. Juozaitis A, Willeke K, Grinshpun SA, Donnelly J. 1994. Impaction onto a glass slide
577 or agar versus impingement into a liquid for the collection and recovery of airborne
578 microorganisms. *Appl Environ Microbiol* 60:861-870.
- 579 41. Cox CS. 1966a. The survival of *Escherichia coli* sprayed into air and into in nitrogen
580 from distilled water and from solutions of protecting agents, as a function of relative
581 humidity. *J Gen Microbiol* 43:383-399.
- 582 42. Marthi B, Lighthart B. 1990. Effects of betaine on enumeration of airborne bacteria.
583 *Appl Environ Microbiol* 56:1286-1289.
- 584 43. Marthi B, Fieland VP, Walter M, Seidler RJ. 1991. Resuscitation effects of catalase
585 on airborne bacteria. *Appl Environ Microbiol* 57:2775-2776.

DSTL/JA100134

- 586 44. Webb, SJ. 1969. The effects of oxygen on the possible repair of dehydration
587 damage by *Escherichia coli*. J Gen Microbiol 58:317-326.
- 588 45. Cox CS. 1968a. The aerosol survival of *Escherichia coli* B in nitrogen, argon and
589 helium atmospheres and the influence of relative humidity. J Gen Microbiol 50:139-
590 147.
- 591 46. Webb SJ, Tai CC. 1968. Lethal and mutagenic action of 3200-4000 Å light. Can J
592 Microbiol 14:727-735.
- 593 47. Xu Z, Wei K, Wu Y, Shen F, Chen Q, Li M, Yao M. 2013. Enhancing bioaerosol
594 sampling by Andersen impactors using mineral-oil-spread agar plate. PLOS One
595 8:e56896
- 596 48. Li C-S, Lin Y-C. 2001. Storage effects on bacterial concentration: determination of
597 impinger and filter samples. Sci Total Environ 278:231-237.
- 598 49. Dybwad M, Skogan G, Blatny JM. 20154. Comparative testing and evaluation of nine
599 different air samplers: end-to-end sampling efficiencies as specific performance
600 measurements for bioaerosols applications. Aero Sci Technol 48:282-295.
- 601 50. Stewart SL, Grinshpun SA, Willeke K, Terzieva S, Ulevicius V & Donnelly J. 1995.
602 Effect of impact stress on microbial recovery on an agar surface. Appl Environ
603 Microbiol 61:1232-1239.
- 604 51. Thomas P, Sekhar AC, Mujawar MM. 2012. Nonrecovery of varying proportions of
605 viable bacteria during spread plating governed by the extent of spreader usage and
606 proposal for an alternate spotting-spreading approach to maximize the CFU. J Appl
607 Microbiol 113:139-150.

DSTL/JA100134

- 608 52. Thomas P, Mujawar MM, Sekhar AC & Upreti R. 2014a. Physical impact injury
609 effects on bacterial cells during spread plating influenced by cell characteristics of
610 the organisms. J Appl Microbiol 116:911-922.
- 611 53. Thomas P, Sekhar AC, Mujawar MM. 2014b. Vulnerability of *Bacillus* spores and of
612 related genera to physical impact injury with particular reference to spread-
613 plating. J Appl Microbiol 117:1358-1372.
- 614 54. Allegra S, Leclerc L, Massard PA, Girardot F, Riffard S, Pourchez S. 2016.
615 Characterization of aerosols containing *Legionella* generated upon nebulization. Sci
616 Rep doi:10.1038/srep33998
- 617 55. Bourouiba L, Dehandschoewercker E, Bush JWM. 2014. Violent expiratory events:
618 on coughing and sneezing. J Fluid Mech 745:537-563.
- 619 56. Scharfman BE, Techet AH, Bush JWM, Bourouiba L. 2016. Visualization of sneeze
620 ejecta: steps of fluid fragmentation leading to respiratory droplets. Exp Fluids
621 57:doi:10.1007/s00348-015-2078-4.
- 622 57. Lai SK, Wang Y-Y, Wirtz D, Hanes J. 2009. Micro- and macrorheology of mucus.
623 Adv Drug Deliv Rev 61:86-100.
- 624 58. Hersen G, Moularat S, Robine E, Ghin E, Corbet S, Vabret A. 2008. Impact of health
625 on particle size of exhaled respiratory aerosols: case-control study. Clean 36:572-
626 577.
- 627 59. Clegg SL, Seinfeld JH, Brimblecombe P. 2001. Thermodynamic modelling of
628 aqueous aerosols containing electrolytes and dissolved organic compounds. J
629 Aerosol Sci 32:713-738.
- 630 60. Power RM, Simpson SH, Reid JP, Hudson AJ. 2013. The transition from liquid to
631 solid-like behaviour in ultrahigh viscosity aerosol particles. Chem Sci 4:2597-2604.

DSTL/JA100134

- 632 61. Elad D, Wolf M, Keck T. 2008. Air-conditioning in the human nasal cavity. *Resp*
633 *Physiol Neurobiol* 163:121-127.
- 634 62. Clegg SL, Seinfeld JH, Edney EO. 2003. Thermodynamic modelling of aqueous
635 aerosols containing electrolytes and dissolved organic compounds. II. An extended
636 Zdanovskii-Stokes-Robinson approach. *J Aero Sci* 34:667-690.
- 637 63. Davies JF, Haddrell AE, Rickards AMJ, Reid JP. 2013. Simultaneous Analysis of the
638 Equilibrium Hygroscopicity and Water Transport Kinetics of Liquid Aerosol. *Anal*
639 *Chem* 85:5819–5826.
- 640 64. Davies JF, Miles REH, Haddrell AE, Reid JP. 2014. Temperature dependence of the
641 vapor pressure and evaporation coefficient of supercooled water, *J Geophys Res -*
642 *Atmos* 119:931–940.
- 643 65. Potts M. 1994. Desiccation tolerance of prokaryotes. *Microbiol Rev* 58:755-805.
- 644 66. Burgess CM, Gianotti A, Gruzdev N, Holah J, Kochel S, Lehner A, Margas E,
645 Schmitz Esser S, Sela S, Tresse O. 2016. The response of foodborne pathogens to
646 osmotic and desiccation stresses in the food chain. *Int J Food Microbiol* 221:37-53.
- 647 67. Johnson DL, Pearce TA, Esmen NA. 1999. The effect of phosphate buffer on
648 aerosol size distribution of nebulized *Bacillus subtilis* and *Pseudomonas fluorescens*
649 bacteria. *Aero Sci Tech* 30:202-210.
- 650 68. Carslaw K S, Clegg SL, Brimblecombe P. 1995. A thermodynamic model of the
651 system HCl-HNO₃-H₂SO₄-H₂O, including solubilities of HBr, from less than 200 to
652 328 K. *J Phys Chem* 99:11557-11574.
- 653 69. Clegg SL, Brimblecombe P. 1995a. A generalized multicomponent thermodynamic
654 model applied to the (NH₄)₂SO₄-H₂SO₄-H₂O system to high supersaturation and low
655 relative humidity at 298.15 K. *J Aero Sci* 26:19-38.

DSTL/JA100134

- 656 70. Clegg SL, Brimblecombe P. 1995b. Application of a multicomponent thermodynamic
657 model to activities and thermal properties of 0-40 mol Kg⁻¹ aqueous sulfuric acid
658 from less than 200-K to 328-K. J Chem Eng Data 40:43-64.
- 659 71. Clegg SL, Ho SS, Chan CK, Brimblecombe P. 1995. Thermodynamic properties of
660 aqueous (NH₄)₂SO₄ to high supersaturation as a function of Ttemperature. J Chem
661 Eng Data 40:1079-1090.
- 662 72. Bowman FM, Melton JA. 2004. Effect of activity coefficient models on predictions of
663 secondary organic aerosol partitioning. J Aerosol Sci 35:1415-1438.
- 664 73. Haddrell AE, Hargreaves G, Davies JF, Reid JP. 2013. Control over hygroscopic
665 growth of saline aqueous aerosol using Pluronic polymer additives. Int J Pharm
666 443:183-192.
- 667 74. Brown AD. 1953. The survival of airborne microorganisms II. Experiments with
668 *Escherichia coli* near 0°C. Aust J Biol Sci 6:470-480.
- 669 75. Won WD, Ross, H. 1966. Effect of diluent and relative humidity on apparent viability
670 of airborne *Pasteurella pestis*. Appl Microbiol 14:742-745.
- 671 76. Cox CS. 1971. Aerosol survival of *Pasteurella tularensis* disseminated from the wet
672 and dry states. Appl Microbiol 21:482-486.
- 673 77. Faith SA, Smith LP, Swatland AS, Reed DS. 2012. Growth conditions and
674 environmental factors impact aerosolization but not virulence of *Francisella*
675 *tularensis* infection in mice. Front Cell Infect Microbiol 2:1-10. doi:
676 10.3389/fcimb.2012.00126.
- 677 78. Handley BA, Webster AJF. 1995. Some factors affecting the airborne survival of
678 bacteria outdoors. J Appl Bacteriol 79:368-378.

DSTL/JA100134

- 679 79. Zuo Z, Kuehn TH, Bekele AZ, Mor SK, Verma H, Goyal SM, Raynor PC. 2014.
680 Survival of airborne MS2 bacteriophage generated from human saliva, artificial
681 saliva, and cell culture medium. *Appl Environ Microbiol* 80:2796-2803.
- 682 80. Ijaz MK, Sattar SA, Johnson-Lussenburg CM, Springthorpe VS, Nair RC. 1985.
683 Effect of relative humidity, atmospheric temperature, and suspending medium on the
684 airborne survival of human rotavirus. *Can J Microbiol* 31:681-685.
- 685 81. Lever MS, Williams A, Bennett AM. 2000. Survival of mycobacterial species in
686 aerosols generated from artificial saliva. *Lett Appl Microbiol* 31:238-241.
- 687 82. Donaldson AI. 1972. The influence of relative humidity on the aerosol stability of
688 different strains of foot-and-mouth disease virus suspended in saliva. *J Gen Virol*
689 15:25-33.
- 690 83. Barlow DF, Donaldson AI. 1973. Comparison of the aerosol stabilities of foot-and-
691 mouth disease virus suspended in cell culture fluid or natural fluids. *J Gen Virol*
692 20:311-318.
- 693 84. Liao CM, Luo WC, Chen SC, Chen JW, Liang HM. 2004. Temporal/seasonal
694 variations of size-dependent airborne fungi indoor/outdoor relationships for a wind-
695 induced naturally ventilated airspace. *Atmos Environ* 38:4415-4419.
- 696 85. Liao CM, Luo WC. 2005. Use of temporal/seasonal- and size-dependent bioaerosol
697 data to characterize the contribution of outdoor fungi to residential exposures. *Sci*
698 *Total Environ* 347:78-97.
- 699 86. Fredenslund A, Jones RL, Prausnitz JM. 1975. Group-contribution estimation of activity
700 coefficients in nonideal liquid mixtures. *AIChE J* 21:1086-1099.

DSTL/JA100134

- 701 87. Peng C, Chan MN, Chan CK. 2001. The hygroscopic properties of dicarboxylic and
702 multifunctional acids: Measurements and UNIFAC predictions. *Environ Sci Technol* 35:
703 4495–4501.
- 704 88. Yttri KE, Dye C, Kiss G. 2007. Ambient aerosol concentrations of sugars and sugar-
705 alcohols at four different sites in Norway. *Atmos Chem Phys* 7, 4267-4279.
- 706 89. Ariya PA, Sun J, Eltouny NA, Hudson ED, Hayes CT, Kos G. 2009. Physical and
707 chemical characterization of bioaerosols – Implications for nucleation processes, *Int*
708 *Rev Phys Chem*, 28:1, 1-32.
- 709 90. Goldberg LJ, Watkins HM, Boerke EE, Chatigny MA. 1958. The use of a rotating
710 drum for the study of aerosols over extended periods of time. *Am J Hyg* 68:85-93.
- 711 91. Gruel RL, Reid CR, Allemann RT. 1987. The optimum rate of drum rotation for
712 aerosol aging. *J Aero Sci* 18:17-22.
- 713 92. Asgharian B, Moss OR. 1992. Particle suspension in a rotating drum chamber when
714 the influence and gravity are both significant. *Aero Sci Tech* 17:263-277.
- 715 93. Krumins V, Son E-K, Mainelis G, Fennell DE. 2008. Retention of inactivated
716 bioaerosols and ethane in a rotating bioreactor constructed for bioaerosols activity
717 studies. *Clean* 36:593-600.
- 718 94. Verrault D, Duchaine C, Marcoux-Voiselle M, Turgeon N, Roy CJ. 2014. Design of
719 an environmentally controlled rotating chamber for bioaerosols aging studies. *Inhal*
720 *Toxicol* 26:554-558.
- 721 95. Druett HA. 1971. A safe method of exposing microthreads in the open air. *J Appl*
722 *Microbiol* 34:655-658.

DSTL/JA100134

- 723 96. May KR, Druett HA. 1968. A microthread technique for studying the viability of
724 microbes in a simulated airborne state. *J Gen Microbiol* 51:353-366.
- 725 97. Handley BA, Roe JM. 1994. An alternative microthread for the study of airborne
726 survival of bacteria outdoors. *J Appl Bacteriol* 77:504-508.
- 727 98. Smither SJ, Piercy TJ, Eastaugh L, Steward JA, Lever MS. 2011. An alternative
728 method of measuring aerosol survival using spiders' webs and its use for filoviruses.
729 *J Virol Meth* 177:123-127.
- 730 99. Adams DJ, Spendlove JC, Spendlove RS, Barnett BB. 1982. Aerosol stability of
731 infectious and potentially infectious reovirus particles. *Appl Environ Microbiol* 44:903-
732 908.
- 733 100. Sattar SA, Ijaz MK, Johnson-Lussenburg CM, Springthorpe CS. 1984. Effect of
734 relative humidity on the airborne survival of rotavirus SA11. *Appl Environ Microbiol*
735 47:879-881.
- 736 101. Deguillaume L, Leriche M, Amato P, Arita PA, Delort A-M, Pöschl U, Chaumerliac
737 N, Bauer H, Flossmann AI, Morris CE. 2008. Microbiology and atmospheric
738 processes: chemical interactions of primary biological aerosols. *Biogeosciences*
739 5:1073-1084.
- 740 102. Estillore AD, Trueblood JV, Grassian VH. 2016. Atmospheric chemistry of
741 bioaerosols: heterogeneous and multiphase reactions with atmospheric oxidants and
742 other trace gases. *Chem Sci* 7:6604-6616.
- 743 103. Cox CS. 1989. Airborne bacteria and viruses. *Sci Prog Oxford* 73:469-500.
- 744 104. Einarsson H, Snygg BG, Eriksson G. 1983. Inhibition of bacterial growth by
745 maillard reaction products. *J Agric Food Chem* 31:1043-1047.

DSTL/JA100134

- 746 105. Hess GE. 1965. Effects of oxygen on aerosolized *Serratia marcescens*. Appl
747 Microbiol 13:781-787.
- 748 106. Benbough JE. 1967. Death mechanisms in airborne *Escherichia coli*. J Gen
749 Microbiol 47:325-333.
- 750 107. Cox CS, Baldwin F. 1967. The toxic effect of oxygen upon the aerosol survival of
751 *Escherichia coli* B. J Gen Microbiol 49:115-117.
- 752 108. Hawkins LN, Lemire AN, Galloway MM, Corrigan AL, Turley JJ., Espelien BM, De
753 Haan DO. 2016. Maillard chemistry in clouds and aqueous aerosol as a source of
754 atmospheric humic-like substances, Environ Sci Technol 50:7443–7452.
- 755 109. Santarpia JL, Pan Y-L, Hill SC, Baker N, Cottrell B, McKee L, Ratnesar-Shumate
756 S, Pinnick RG. 2012. Changes in fluorescence spectra of bioaerosols exposed to
757 ozone in a laboratory reaction chamber to simulate atmospheric aging. Optics
758 Express 20:29867-29881.
- 759 110. Pan Y-L, Santarpia JL, Ratnesar-Shumate S, Corson E, Eshbaugh J, Hill SC,
760 Williamson CC, Coleman M, Bare C, Kinahan S. 2014. Effects of ozone and relative
761 humidity on fluorescence spectra of octapeptide bioaerosol particles. J Quant Spect
762 Rad Trans 538-550.
- 763 111. Ratnesar-Shumate S, Pan Y-L, Hill SC, Kinahan S, Corson E, Eshbaugh J,
764 Santarpia JL. 2015. Fluorescence spectra and biological activity of aerosolized
765 *Bacillus* spores and MS2 bacteriophage exposed to ozone at different relative
766 humidities in a rotating drum. J Quant Spect Rad Trans 153:13-28.
- 767 112. Cox CS. 1970. Aerosol survival of *Escherichia coli* B disseminated from the dry
768 state. Appl Microbiol 19:604-607.

DSTL/JA100134

- 769 113. Cox CS. 1966b. The survival of *Escherichia coli* in nitrogen atmospheres under
770 changing conditions of relative humidity. J Gen Microbiol 45:283-288.
- 771 114. Yang W, Marr, LC. 2012. Mechanisms by which ambient humidity may affect
772 viruses in aerosols. Appl Environ Microbiol 78:6781-6788.
- 773 115. Beebe JM. 1959. Stability of disseminated aerosols of *Pasteurella tularensis*
774 subjected to simulated solar radiations at various humidities. J Bacteriol 78:18-24.
- 775 116. Tong Y, Lighthart B. 1997. Solar radiation has a lethal effect on natural
776 populations of culturable outdoor atmospheric bacteria. Atmos Environ 31:897-900.
- 777 117. Tong Y, Lighthart B. 1998. Effect of simulated solar radiation on mixed outdoor
778 atmospheric bacterial populations. FEMS Microbiol Ecol 26:311-316.
- 779 118. Berendt RF, Dorsey EL. 1971. Effect of simulated solar radiation and sodium
780 fluorescein on the recovery of Venezuelan Equine Encephalomyelitis virus from
781 aerosols. Appl Microbiol 21:447-450
- 782 119. Gray LJ, Beer J, Geller M, Haigh JD, Lockwood M, Matthes K, Cubasch U,
783 Fleitmann D, Harrison G, Hood L, Luterbacher J, Meehl GA, Shindell D, van Geel B,
784 White W. 2010. Solar influences on climate. Rev Geophys 48:1-53.
- 785 120. Rastogi RP, Richa, Kumar A, Tyagi MB, Sinha RP. 2010. Molecular mechanisms
786 of ultraviolet radiation-induced DNA damage and repair. J Nucleic Acids
787 doi:10.4061/2010/592980.
- 788 121. Santos AL, Gomes NC, Henriques I, Almeida A, Correia A, Cunha A. 2012.
789 Contribution of reactive oxygen species to UV-B induced damage in bacteria. J
790 Photochem Photobiol B 117:40-46.

DSTL/JA100134

- 791 122. de Mik G, de Groot I. 1977. Mechanisms of inactivation of bacteriophage ϕ X174
792 and its DNA in aerosols by ozone and ozonized cyclohexene. J Hyg – Camb 78:199-
793 211.
- 794 123. Dark FA, Nash T. 1970. Comparative toxicity of various ozonized olefins to
795 bacteria suspended in air. J Hyg – Camb 68:245-252.
- 796 124. Hood AM. 1971. An indoor system for the study of biological aerosols in open air
797 conditions. J Hyg – Camb 69:607-617.
- 798 125. Donaldson AI, Ferris NP. 1975. The survival of foot-and-mouth disease virus in
799 open air conditions. J Hyg (Lond) 74:409-416.
- 800 126. Lighthart B. 1973. Survival of airborne bacteria in a high urban concentration of
801 carbon monoxide. Appl Microbiol 25:86-91.
- 802 127. Lighthart B, Hiatt VE, Rossano AT. 1971. The survival of airborne *Serratia*
803 *marcescens* in urban concentrations of sulphur dioxide. J Air Pollut Cont Assoc
804 21:639-642.
- 805 128. Ehrlich R, Miller S. 1972. Effect of NO₂ on airborne Venezuelan Equine
806 Encephalomyelitis virus. Appl Microbiol 23:481-484.
- 807 129. Hood AM. 2009. The effect of open-air factors on the virulence and viability of
808 airborne *Francisella tularensis*. Epidemiol Infect 137:753-761.
- 809 130. Benbough JE, Hood AM. 1971. Viricidal activity of open air. J Hyg Camb 69:619-
810 626.
- 811 131. de Mik G, de Groot I. 1978. Breaks induced in the deoxyribonucleic acid of
812 aerosolized *Escherichia coli* by ozonized cyclohexene. Appl Environ Microbiol 35:6-
813 10.

DSTL/JA100134

- 814 132. Anderson JD. 1966. Biochemical studies of lethal processes in aerosols of
815 *Escherichia coli*. J Gen Microbiol 45:303-313.
- 816 133. Cox CS. 1969. The cause of loss of viability of airborne *Escherichia coli* K12. J
817 Gen Microbiol 57:77-80.
- 818 134. Csonka LN. 1989. Physiological and genetic responses of bacteria to osmotic
819 stress. Microbiol Rev 53:121-147.
- 820 135. Sleator RD, Hill C. 2002. Bacterial osmoadaptation: the role of osmolytes in
821 bacterial stress and virulence. FEMS Microbiol Rev 26:49-71.
- 822 136. Wood JM. 2015. Bacterial responses to osmotic challenges. J Gen Physiol
823 145:381-388.
- 824 137. Anderson JD, Dark FA, Peto S. 1968. Studies on the effects of aerosolization on
825 the rates of efflux of ions from populations of *Escherichia coli* strain B. J Gen
826 Microbiol 46:95-105.
- 827 138. Gruzdev N, McClelland M, Porwollik S, Ofaim S, Pinto R, Saldinger-Sela S. 2012.
828 Global transcriptional analysis of dehydrated *Salmonella enterica* serovar
829 Typhimurium. Appl Environ Microbiol 78:7866-7875.
- 830 139. Kocharunchitt C, King T, Gobius K, Bowman JP, Ross T. 2014. Global genome
831 response of *Escherichia coli* O157:H7 Sakai during dynamic changes in growth
832 kinetics induced by an abrupt downshift in water activity. PLOS One 9:e90422
- 833 140. Fung KH, Tang IN. 2000. Chemical characterization of aerosol particles by laser
834 raman spectroscopy, p 177-195. In Spurny KR (ed), Aerosol Chemical Processes in
835 the Environment. CRC Press LLC, Florida.

DSTL/JA100134

- 836 141. Rubel GO. 2000. Novel applications of the electrodynamic levitator for the study
837 of aerosol chemical processes, p 197-211. *In* Spurny KR (ed), Aerosol Chemical
838 Processes in the Environment. CRC Press LLC, Florida.
- 839 142. Reid JP, Meresman H, Mitchem L, Symes R. 2007. Spectroscopic studies of the
840 size and composition of single aerosol droplets. *Int Rev Phys Chem* 26:139–192.
- 841 143. Reid JP. 2009. Particle levitation and laboratory scattering. *J Quant Spectrosc*
842 *Radiat Transf* 110:1293–1306.
- 843 144. Ault AP, Zhao D, Ebben CJ, Tauber MJ, Geiger FM, Prather KA, Grassian VH.
844 2013. Raman microspectroscopy and vibrational sum frequency generation
845 spectroscopy as probes of the bulk and surface compositions of size-resolved sea
846 spray aerosol particles. *Phys Chem Chem Phys* 15:6206-6214.
- 847 145. Craig RL, Bondy AL, Ault AP. 2015. Surface enhanced raman spectroscopy
848 enables observations of previously undetectable secondary organic aerosol
849 components at the individual particle level. *Anal Chem* 87:7510-7514.
- 850 146. Xie C, Li Y-Q. 2003. Confocal micro-Raman spectroscopy of single biological
851 cells using optical trapping and shifted excitation difference techniques. *J Appl Phys*
852 93:2982-2986.
- 853 147. Kalasinsky KS, Hadfield T, Shea AA, Kalasinsky VF, Nelson MP, Neiss J, Drauch
854 AJ, Vanni GS, Treado PJ. 2007. Raman chemical imaging spectroscopy reagentless
855 detection and identification of pathogens: signature development and evaluation.
856 *Anal Chem* 79:2658-2673.

DSTL/JA100134

- 857 148. Meisel S, Stöckel S, Elschner M, Melzer F, Rösch P, Popp J. 2012. Raman
858 spectroscopy as a potential tool for detection of *Brucella* spp. in milk. Appl Environ
859 Microbiol 78:5575-5583.
- 860 149. Zu TNK, Athamneh AIM, Wallace RS, Collakova E, Senger RS. 2014. Near real-
861 time analysis of the phenotypic response of *Escherichia coli* to 1-butanol exposure
862 using Raman spectroscopy. J Bacteriol 196:3983-3991.
- 863 150. Sengupta A, Laucks ML, Dildine N, Drapala E, Davis EJ. 2005. Bioaerosol
864 characterization by surface-enhanced Raman spectroscopy (SERS). Aero Sci
865 36:651-664.
- 866 151. Sengupta A, Brar N, Davis EJ. 2007. Bioaerosol detection and characterization
867 by surface-enhanced Raman spectroscopy. J Colloid Interface Sci 309:36-43.
- 868 152. Wang C, Pan Y-L, Hill SC, Redding B. 2015. Photophoretic trapping-Raman
869 spectroscopy for single pollens and fungal spores trapped in air. J Quant Spect Rad
870 Trans 153:4-12.
- 871 153. Cruz-Sanchez TM, Haddrell AE, Hackett TL, Singhera GK, Marchant D, Lekivetz
872 R, Meredith A, Horne D, Knight DA, van Eeden SF, Bai TR, Hegele RG, Dorscheid
873 DR, Agnes GR. 2013. Formation of a stable mimic of ambient particulate matter
874 containing viable infectious respiratory syncytial virus and its dry-deposition directly
875 onto cell cultures. Anal Chem 85:898-906.
- 876 154. Hingston PA, Piercey MJ, Hansen LT. 2015. Genes associated with desiccation
877 and osmotic stress in *Listeria monocytogenes* as revealed by insertional
878 mutagenesis. Appl Environ Microbiol 81:5350-5362.
- 879 155. Wang D, Bodovitz S. 2010. Single cell analysis: the new frontier in 'Omics'.
880 Trends Biotechnol 28:281-290.

DSTL/JA100134

- 881 156. Heinemann M, Zenobi R. 2011. Single cell metabolomics. *Curr Opin Biotechnol*
882 22:26-31.
- 883 157. Blainey PC. 2013. The future is now: single-cell genomics of bacteria and
884 archaea. *FEMS Microbiol Rev* 37:1-29.
- 885 158. Tsioris K, Torres AJ, Douce TB, Love JC. 2014. A new toolbox for assessing
886 single cells. *Ann Rev Chem Biomol Eng* 5:455-477.
- 887 159. Kalisky T, Quake SR. 2011. Single-cell genomics. *Nat Meth* 8:311-314.
- 888 160. Hall RJ, Leblanc-Maridor M, Wang J, Ren X, Moore NE, Brooks CR, Peacey M,
889 Douwes J, McLean D. (2013). Metagenomic detection of viruses in animal
890 slaughterhouses. *PLoS One* 8:e72226.
- 891 161. Be NA, Thissen JB, Fofanov VY, Allen JE, Rojas M, Golovko G, Fofanov Y,
892 Koshinsky H, Jiang CJ. 2015. Metagenomic analysis of the airborne environment in
893 urban spaces. *Microb Ecol* 69:346-355.
- 894 162. May KR. 1973. The Collison nebulizer: description, performance and application.
895 *Aero Sci* 4:235-243.
- 896 163. Cox CS, Goldberg LJ. 1972. Aerosol survival of *Pasteurella tularensis* and the
897 influence of relative humidity. *Appl Microbiol* 23:1-3.
- 898 164. Ehrlich R, Miller S. 1973. Survival of airborne *Pasteurella tularensis* at different
899 atmospheric temperatures. *Appl Microbiol* 25:369-372.
- 900 165. Cox CS, Bondurant MC, Hatch MT. 1971. Effects of oxygen on aerosol survival
901 of radiation sensitive and resistant strains of *Escherichia coli* B. *J Hyg – Camb*
902 69:661-672.

DSTL/JA100134

- 903 166. Woo M-H, Grippin A, Anwar D, Smith T, Wu C-Y, Wander JD. 2012. Effects of
904 relative humidity and spraying medium on UV decontamination of filters loaded with
905 viral aerosols. *Appl Environ Microbiol* 78:5781-5787.
- 906 167. Chen BT, John W. 2001. Instrument Calibration. *In* Baron PA, Willeke K (eds).
907 Aerosol Measurement: Principles, Techniques, and Applications. Wiley, New York.
- 908 168. Young HW, Larson EW, Dominik JW. 1974. Modified spinning top homogeneous
909 spray apparatus for use in experimental respiratory disease studies. *Appl Microbiol*
910 28:929-934.
- 911 169. Thomas RJ, Webber D, Sellors W, Collinge A, Stagg AJ, Bailey SC, Jayasekera
912 PN, Taylor RR, Eley SE, Titball RW. 2008. Characterization and deposition of
913 respirable large- and small-particle bioaerosols. *Appl Environ Microbiol* 74:6437-
914 6443.
- 915 170. Grinshpun SA, Willeke K, Ulevicius V, Juozaitis A, Terzieva S, Donnelly J,
916 Stelma GN, Brenner KP. 1997. Effect of impaction, bounce and reaerosolization on
917 the collection efficiency of impingers. *Aero Sci Technol* 26:326-342.
- 918 171. Li C-S, Hao M-L, Lin W-H, Chang C-W, Wang C-S. 1999. Evaluation of aerosol
919 samplers for bacterial microorganisms. *Aero Sci Technol* 30:100-108.
- 920 172. Lin X, Reponen TA, Willeke K, Grinshpun SA, Foarde KK, Ensor DS. 1999. Long-
921 term sampling of airborne bacteria and fungi into a non-evaporating liquid. *Atmos*
922 *Environ* 33:4291-4298.
- 923 173. Lin X, Reponen T, Willeke K, Wang Z, Grinshpun SA, Trunov M. 2000. Survival
924 of airborne microorganisms during swirling aerosol collection. *Aero Sci Technol*
925 32:184-196.

DSTL/JA100134

- 926 174. Riemenschneider L, Woo W-H, Wu C-Y, Lundgren D, Wander J, Lee J-H & Li H-
927 W. 2009. Characterization of reaerosolization from impingers in an effort to improve
928 airborne virus sampling. *J Appl Microbiol* 108:315-324.
- 929 175. May KR, Harper GJ. 1957. The efficiency of various liquid impinger samplers in
930 bacterial aerosols. *Brit J Industr Med* 14:287-297.
- 931 176. Kesavan J, Schepers D, McFarland AR. 2010. Sampling and retention
932 efficiencies of batch-type liquid-based bioaerosols samplers. *Aero Sci Technol*
933 44:817-829.
- 934 177. Yu KP, Chen Y-P, Gong J-Y, Cen Y-C, Cheng C-C. 2016. Improving the
935 collection efficiency of the liquid impinger for ultrafine particles and viral aerosols by
936 applying granular bed filtration. *J Aero Sci* 101:133-143.
- 937 178. Zheng Y, Yao M. 2017. Liquid impinger Biosampler's performance for size-
938 resolved viable bioaerosol particles. *J Aero Sci* 106:34-42.
- 939 179. Andersen AA. 1958. New sampler for the collection, sizing and enumeration of
940 viable airborne particles. *J Bacteriol* 76:471-484.
- 941 180. May KR. 1975. An "ultimate" cascade impactor for aerosol assessment. *J Aero*
942 *Sci* 6:413-419.
- 943 181. Li C-S, Lin Y-C. 1999. Sampling performance of impactors for bacterial
944 bioaerosols. *Aero Sci Technol* 30:280-287.
- 945 182. Mainelis G, Willeke K, Adhikari A, Reponen T, Grinshpun SA 2002. Design and
946 collection efficiency of a new electrostatic precipitator for bioaerosol collection. *Aero*
947 *Sci Technol* 36:1073-1085.
- 948 183. Cheng YS. 2014. Mechanisms of pharmaceutical aerosol deposition in the
949 respiratory tract. *AAPS Pharm Sci Tech* 15:630-640.

DSTL/JA100134

- 950 184. Rabey F, Janssen RJ, Kelley LM. 1969. Stability of St. Louis Encephalitis virus in
951 the airborne state. *Appl Microbiol* 18:880-882.
- 952 185. Hood AM. 1974. Open-air factor in enclosed systems. *J Hyg* 72:53-60.
- 953 186. Bailey R, Fielding L, Young A, Griffith C. 2007. Effect of ozone and open air
954 factor against aerosolized *Micrococcus luteus*. *J Food Prot* 70: 2679-2773.
- 955 187. Walter MV, Marthi B, Fieland VP, Ganio LM. 1990. Effect of aerosolization on
956 subsequent bacterial survival. *Appl Environ Microbiol* 56:3468-3472.
- 957 188. Marthi B, Fieland VP, Walter M, Seidler RJ. 1990. Survival of bacteria during
958 aerosolization. *Appl Environ Microbiol* 56:3463-3467.
- 959 189. Cox CS. 1968b. The aerosol survival and cause of death of *Escherichia coli* K12.
960 *J Gen Microbiol* 54:169-175.
- 961 190. Hatch MT, Dimmick RL. 1966. Physiological responses of airborne bacteria to
962 shifts in relative humidity. *Bacteriol Rev* 30:597-602.
- 963 191. Ehrlich R, Miller S. 1971. Effect of relative humidity and temperature on airborne
964 Venezuelan Equine Encephalitis virus. *Appl Microbiol* 22:194-199.
- 965 192. Wathes CM, Howard K, Webster AJF. 1986. The survival of *Escherichia coli* in an
966 aerosol at air temperatures of 15 and 30 °C and a range of humidities. *J Hyg Camb*
967 97:489-496.
- 968 193. Thompson K-A, Bennett AM, Walker JT. 2011. Aerosol survival of
969 *Staphylococcus epidermidis*. *J Hosp Infect* 78:216-220.
- 970 194. Ehrlich R, Miller S, Walker RL. 1970. Relationship between atmospheric
971 temperature and survival of airborne bacteria. *Appl Microbiol* 19:245-249.

DSTL/JA100134

- 972 195. Benbough JE. 1969. Factors affecting the toxicity of oxygen towards airborne
973 coliform bacteria. J Gen Microbiol 56:241-250.
- 974 196. Cox CS, Derr JS, Flurie EG, Roderick RC. 1970. Experimental technique for
975 studying aerosols of lyophilized bacteria. Appl Microbiol 20:927-934.
- 976 197. Schlamm NA. 1960. Detection of viability in aged or injured *Pasteurella*
977 *tularensis*. J Bacteriol 91:2180-2184.
- 978 198. Sawyer WD, Jemski JV, Hogge AL, Eigelsbach HT, Wolfe EK, Dangerfield HG,
979 Gochenour WS, Crozier D. 1966. Effect of aerosol age on the infectivity of airborne
980 *Pasteurella tularensis* for *Mucaca mulatta* and man. J Bacteriol 91:2180-2184.
- 981 199. Hood AM. 1961. Infectivity of *Pasteurella tularensis* clouds. J Hyg 59:497-504.
- 982 .
- 983
- 984
- 985
- 986
- 987
- 988
- 989
- 990
- 991

DSTL/JA100134

993 TABLE 1 Methods used to generate and sample microbial aerosols useful for aerosol fate and inhalational infection research.

Mechanism	Apparatus examples	Description	References(s)
Aerosol generation			
Reflux nebulization (1-, 3-, 6-jet versions commonly used)	Collison nebulizer, Wells atomizer, TSI 9302, FK-8 aerosol gun, Aeroneb Lab	<ul style="list-style-type: none"> Refluxing two-fluid atomizer operating by venturi effect and wall impaction. Liquid recirculation occurs every 6 seconds in the 3-jet version (135). Increased jet numbers increase the rate of aerosol generation and recirculation. Reservoir evaporation occurs over time causing concentration effects. Generally used for liquids, although the Wells atomizer was used for dry powders. Particle sizes are small, 0.7–2.2 μm. Forces associated with reflux nebulization can cause deagglomeration of aggregates causing an observed increase in bacterial concentration in the spray suspension. 	14, 16, 20, 23-25, 79, 80, 99, 122, 162-167
Non-reflux nebulization	Single-pass aerosolizer	<ul style="list-style-type: none"> Atomisation as above minus wall impaction and recirculation 	24
Aerosol bubbling	SLAG ^p and variants	<ul style="list-style-type: none"> Liquid dripped onto a membrane is broken into droplets by air flow through the membrane. Droplets burst due to increased pressure gradient between the inside and outside 	16, 24, 26

DSTL/JA100134

		of the device generating small aerosol particles.	
Centrifugal atomization	Spinning top aerosol generator	<ul style="list-style-type: none">Centrifugal forces moves liquid applied to a rotating disc towards the edges producing ligands that break into droplets	168
Flow-focussing	FFAG ^o , C-Flow nebulizer	<ul style="list-style-type: none">Liquid flows through an orifice forming microjets that break-up into particles by aerodynamic suction of an accelerated air stream.Good monodispersity of droplets can be achieved.	20, 24, 169
Aerosol sampling			
Impingement	Impingers ^d (AGI-4, AGI-30, Model 7541 AGI); SKC biosampler	<ul style="list-style-type: none">Aerosol accelerates through critical orifice causing inertial impaction into liquid.Efficiency is affected by physical parameters (e.g. sampling flow rate, nozzle number and angle, distance of nozzle from the liquid, solution type and volume, particle bounce, prolonged sampling time (liquid evaporation, increased damage) and binding of microorganisms to the collection vessel wall.Reaerosolization can occur due to liquid bubbling.Addition of glass beads can increase virus collection efficiencySKC biosampler possesses three angled nozzles creating a gentler swirling	17, 18, 21, 22, 170-178

DSTL/JA100134

		<p>motion of the bioaerosol during collection.</p> <ul style="list-style-type: none"> • AGI-30 impaction velocity reaches $265 \text{ m}\cdot\text{s}^{-1}$; much reduced in other samplers. 	
Impaction	<p>Single or multistage impactors:</p> <p>Andersen, Mercer, Ultimate, MAS-100, Burkard</p>	<ul style="list-style-type: none"> • Operate at constant flow rates, with air flowing through an orifice causing inertial impaction of particles too large to remain entrained in the air flow; size fractionation possible. • Collection can be onto a range of different substrates (e.g. agar plates, gelatin coated slides or filters). • Substrate choice can affect collection efficiency due to effects on microbial viability and particle bounce. • In the Burkard and 6th stage of the Andersen impactors, impaction velocities reach 12 and $24 \text{ m}\cdot\text{s}^{-1}$ respectively. 	21, 22, 47, 179-181
Filtration/impaction	<p>Gelatin filter, nitrocellulose, polycarbonate</p>	<ul style="list-style-type: none"> • Greater physical sampling efficiencies. Biological sampling efficiency may be lower due to sensitivity of the collected microorganisms to air drawn past the filter. • Elution of material from the filter surface (e.g. vortexing, shaking, solution volume and type) can influence efficiency. 	21, 22, 48, 49

DSTL/JA100134

Direct capture	Microthreads	<ul style="list-style-type: none"> • Particles collected onto fine microthreads (e.g. spider silk, glue thread) wound on to a frame. 	78, 96-98, 123-125
Cyclonic separation	NIOSH cyclonic biosampler	<ul style="list-style-type: none"> • Air flow drawn into a cylindrical container is rotated causing larger particles to deposit and collect on the walls by centrifugal forces. 	25, 35
Electrostatic precipitation	Ionizers e.g. AS150; Model 3100 aerosol sampler	<ul style="list-style-type: none"> • Airborne particles electrically charged and subjected to electric field causing gentle deposition velocity onto collection substrate. • Bioefficiency for spores greater than for Gram-negative bacteria. • Impaction velocities reach $0.01 - 1 \text{ m}\cdot\text{s}^{-1}$. 	30, 36, 182
Animal inhalation	Rodent, primates	<ul style="list-style-type: none"> • Aerosol particles regionally deposit due to inertial impaction, sedimentation, diffusion, interception and electrostatic effects in the respiratory tract. • Deposition is a function of airway geometry and particle properties (e.g. size, shape, density, hygroscopicity). 	183

994

995 ^a = Note that the list is merely representative and not exhaustive. Researchers are recommended to conduct rigorous validation of
 996 the aerosol experimental system for each individual micro-organism tested; ^b = sparging liquid aerosol generator; ^c = flow focussing
 997 aerosol generator; ^d = all-glass impinger

DSTL/JA100134

998 TABLE 2 Examples of experimental techniques used to study fate of microorganisms
999 in aerosol

Device	Mechanism	Aerosol state ^a	Outdoor use?	References(s)
Rotating drum	Rotational speed of drum	Dynamic	N	34, 82, 83,
	prevents aerosol from settling			93, 94, 99,
	for period of time dependent			111, 126,
	on particle size			163, 184
Microthread	Aerosol captured on spider	Captured	Y	78, 96-98, 123-125, 130
	microthreads or glue fibres			
	wound around a metal frame			
	that can be slotted into an exposure apparatus.			
Sphere	Steel sphere with mixing fans	Dynamic	N	124, 185
Aerosol chamber	Large chambers with mixing fans	Dynamic	N	186
Greenhouse	No mixing fan	Dynamic	Y	187,188

1000 ^a = Dynamic refers to particles maintained as a buoyant aerosol, whilst captured

1001 refers to aerosol particles immobilised on a substrate.

1002

1003

DSTL/JA100134

1004 TABLE 3 Atmospheric, environmental and microbial factors affecting survival and
1005 infectivity in airborne microorganisms

Factor	Description	References(s) ^a
Relative humidity (RH)	Generally studies range from 20 to 90% RH	41, 45, 76, 80, 82, 99, 100, 113, 115, 163, 184, 189-193
Temperature	Wide ranges studied from sub-zero to 50 °C	80, 164, 191, 192, 194
Solar radiation	Variability in spectra examined but inclusive of UV-A and UV-B wavelengths	46, 78, 115-118, 188
Oxygen	Generation of ROS ^b during aerosol transport	44, 105-107, 165, 195
Ozone	Reactive with pollutant gases and pinenes	122, 186
Pollutant gases 'Open air factor'	CO, SO ₂ , NO ₂ , ethene, cyclohexene SOAs ^c (e.g. alkenes, turpenes ^d)	31, 93, 122-126, 127-131, 185
Wet / dry preparation	Droplets or dried particles	76, 112, 163, 189, 196
Growth phase	Exponential, stationary	31, 165
Particle size	Microbial aggregates have greater survival than single microorganisms	31, 78, 130, 195
Aerosol age	Infectivity decreased prior to culturability with extended time in aerosol	197-199

DSTL/JA100134

- 1006 ^a = reference list is reflective and not exhaustive; ^b = reactive oxygen species; ^c =
1007 secondary organic aerosol; ^d = turpenes are volatile cyclic unsaturated hydrocarbon
1008 molecules released by plants